



<http://ijgt.ui.ac.ir>



www.ui.ac.ir

ON CO-MAXIMAL SUBGROUP GRAPH OF \mathbb{Z}_n

MANIDEEPA SAHA, SUCHARITA BISWAS AND ANGSUMAN DAS*

ABSTRACT. The co-maximal subgroup graph $\Gamma(G)$ of a group G is a graph whose vertices are non-trivial proper subgroups of G and two vertices H and K are adjacent if $HK = G$. In this paper, we study and characterize various properties like diameter, domination number, perfectness, hamiltonicity, etc. of $\Gamma(\mathbb{Z}_n)$.

1. Introduction

Over the last two decades, various graphs has been defined on groups. For a comprehensive survey on this, please refer to [2]. One of such graphs, the *co-maximal subgroup graph* of a group G was introduced by Akbari *et.al.* [1] in 2017. More results on comaximal subgroup graph can be found in [3] and [4]. In this paper, we study various properties of co-maximal subgroup graph of \mathbb{Z}_n . In particular, we characterize the value of n for which the graph is hamiltonian, eulerian, perfect etc.

We first recall a few definitions and results from [1] and [3].

Definition 1.1. *Let G be a group and S be the collection of all non-trivial proper subgroups of G . The co-maximal subgroup graph $\Gamma(G)$ of a group G is defined to be a graph with S as the set of vertices and two distinct vertices H and K are adjacent if and only if $HK = G$.*

Theorem 1.2. [1], [3] *Let G be a finite nilpotent group.*

Communicated by Attila Maroti.

MSC(2010): Primary: 05C25; Secondary: 05C17, 05C45.

Keywords: hamiltonicity, isomorphism, perfect graph.

Received: 06 August 2021, Accepted: 29 September 2021.

Article Type: Research Paper.

*Corresponding author.

<http://dx.doi.org/10.22108/IJGT.2021.129788.1732>

- $\Gamma(G)$ is connected apart from possible isolated vertices, which corresponds to the non-trivial subgroups of the Frattini subgroup of G .
- Denote by $\Gamma^*(G)$, the graph obtained from $\Gamma(G)$ by removing the isolated vertices, if any. Then $\text{diam}(\Gamma^*(G)) \leq 3$.
- $\Gamma^*(G)$ is weakly perfect, i.e., the clique number and the chromatic number of $\Gamma^*(G)$ are equal.

As cyclic groups are nilpotent, all the above results hold for $\Gamma(\mathbb{Z}_n)$. We start with some basic properties of $\Gamma(\mathbb{Z}_n)$ and $\Gamma^*(\mathbb{Z}_n)$. As for any cyclic p -group G , $\Gamma(G)$ is empty, throughout the paper, we consider $\Gamma(\mathbb{Z}_n)$ where n is not a prime power.

2. Basic Properties

In this section, we study some basic properties of $\Gamma(\mathbb{Z}_n)$ and $\Gamma^*(\mathbb{Z}_n)$ like connectedness, degree, diameter etc.

Lemma 2.1. Let $H = \langle x \rangle$ and $K = \langle y \rangle$ be two subgroups of \mathbb{Z}_n where x, y divide n . Then $H \sim K$ in $\Gamma(\mathbb{Z}_n)$ if and only if $\gcd(x, y) = 1$.

Proof. It follows from Bezout's Theorem and the observation that $HK = \{sx + ty : s, t \in \mathbb{Z}\}$. \square

Theorem 2.2. Let $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$, where p_i 's are distinct primes and $\alpha_i \geq 1$. Let $H = \langle p_1^{\beta_1} p_2^{\beta_2} \cdots p_k^{\beta_k} \rangle$ be a subgroup of \mathbb{Z}_n , where $\beta_i \leq \alpha_i$. Then degree of H in $\Gamma(\mathbb{Z}_n)$ is

$$\deg(H) = \begin{cases} 0, & \text{if } \beta_i \neq 0, \forall i, \\ \prod_{j:\beta_j=0} (\alpha_j + 1) - 1, & \text{otherwise.} \end{cases}$$

Proof. Follows from Lemma 2.1. \square

Corollary 2.3. Let $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$, where p_i 's are distinct primes and $\alpha_i \geq 1$. Then $\Gamma^*(\mathbb{Z}_n)$ is Eulerian if and only if n is a perfect square.

Proof. If n is a perfect square, then each α_i is even and by Theorem 2.2, degree of every vertex of $\Gamma^*(\mathbb{Z}_n)$ is even and hence $\Gamma^*(\mathbb{Z}_n)$ is Eulerian. If n is not a perfect square, then there exists i such that α_i is odd. Let $H = \langle p_1^{\alpha_1} \cdots p_{i-1}^{\alpha_{i-1}} p_{i+1}^{\alpha_{i+1}} \cdots p_k^{\alpha_k} \rangle$. Then by Theorem 2.2, $\deg(H) = \alpha_i$, which is odd. Thus $\Gamma^*(\mathbb{Z}_n)$ is not Eulerian. \square

Theorem 2.4. Let $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$, where p_i 's are distinct primes and $\alpha_i \geq 1$. Then $\Gamma(\mathbb{Z}_n)$ has exactly $\alpha_1 \alpha_2 \cdots \alpha_k - 1$ isolated vertices.

Proof. Since G is a cyclic non p -group of order $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$, $G \cong \mathbb{Z}_n$. By Lemma 2.1, $H = \langle p_1 p_2 \cdots p_k \rangle$ is an isolated vertex in $\Gamma(G)$. Similarly, if x is a multiple of $p_1 p_2 \cdots p_k$ which divides n , then $\langle x \rangle$ is an isolated vertex in $\Gamma(G)$.

Let $A = \langle a \rangle$ with $a|n$ be a subgroup of G such that A is an isolated vertex in $\Gamma(G)$. As G has a unique subgroup of order corresponding to each factor of n and for any non-trivial proper subgroup H of G , we deduce $A \not\sim H$ in $\Gamma(G)$, we have $\gcd(a, m) \neq 1$ for any factor m of $|G| = n$. Thus $p_i|a$ for all i , i.e., a is a multiple of $p_1p_2 \cdots p_k$ which divides n .

Hence the number of isolated vertices in $\Gamma(G)$ is $\alpha_1\alpha_2 \cdots \alpha_k - 1$. □

Corollary 2.5. $\Gamma(\mathbb{Z}_n)$ is connected if and only if n is square-free.

Proof. Let $n = p_1^{\alpha_1}p_2^{\alpha_2} \cdots p_k^{\alpha_k}$. Then the corollary follows from the fact that $\alpha_1\alpha_2 \cdots \alpha_k - 1 = 0$ if and only if n is square-free. □

Theorem 2.6. Let G be a cyclic non p -group of order $n = p_1^{\alpha_1}p_2^{\alpha_2} \cdots p_k^{\alpha_k}$, where p_i 's are distinct primes and $\alpha_i \geq 1$. Then $\text{diam}(\Gamma^*(G)) = \begin{cases} 2, & \text{if } k = 2, \\ 3, & \text{if } k \geq 3. \end{cases}$

Proof. It is clear that the number of maximal subgroups of G is k . If $k = 2$, then the vertices of $\Gamma^*(G)$ are $\langle p_1 \rangle, \langle p_1^2 \rangle, \dots, \langle p_1^{\alpha_1} \rangle, \langle p_2 \rangle, \langle p_2^2 \rangle, \dots, \langle p_2^{\alpha_2} \rangle$ and any two non-adjacent vertices always have a common neighbour either $\langle p_1 \rangle$ or $\langle p_2 \rangle$. Hence its diameter is 2.

If $k \geq 3$, then $\langle p_1p_2 \cdots p_{k-1} \rangle$ and $\langle p_2p_3 \cdots p_k \rangle$ are non-adjacent vertices in $\Gamma^*(G)$ and they do not have any common neighbour. Thus their distance is greater than 2. Now, as \mathbb{Z}_n is nilpotent, $\text{diam}(\Gamma^*(G)) = 3$. □

Theorem 2.7. Let G be a cyclic non p -group of order $n = p_1^{\alpha_1}p_2^{\alpha_2} \cdots p_k^{\alpha_k}$, where p_i 's are distinct primes and $\alpha_i \geq 1$. Then $\Gamma(G)$ has pendant vertices if and only if $\alpha_i = 1$ for some i .

Proof. Let G be a cyclic group of order $n = p_1^{\alpha_1}p_2^{\alpha_2} \cdots p_k^{\alpha_k}$, where at least one $\alpha_i = 1$, say $\alpha_1 = 1$, i.e., $n = p_1p_2^{\alpha_2} \cdots p_k^{\alpha_k}$. Then $\langle p_2p_3 \cdots p_k \rangle$ is a pendant vertex in $\Gamma(G)$, which is adjacent to $\langle p_1 \rangle$.

Conversely, let G be a cyclic group of order $n = p_1^{\alpha_1}p_2^{\alpha_2} \cdots p_k^{\alpha_k}$ such that $\Gamma(G)$ has at least one pendant vertex. If possible, let $\alpha_i \geq 2$ for all i . Let $H = \langle m \rangle$ be a pendant vertex in $\Gamma(G)$ where $m|n$. If $p_i|m$ for all i , then H is an isolated vertex, a contradiction. Thus, m misses at least one prime factor. Let $m = p_2^{\beta_2} \cdots p_k^{\beta_k}$ where $0 \leq \beta_i \leq \alpha_i$. But this implies that H is adjacent to the vertices $\langle p_1 \rangle, \langle p_1^2 \rangle, \dots, \langle p_1^{\alpha_1} \rangle$. As $\alpha_1 \geq 2$, H can not be a pendant vertex. Thus, at least some α_i must be 1. □

3. Hamiltonicity, Perfectness and Dominating Sets

In this section, we characterize the values of n for which $\Gamma^*(\mathbb{Z}_n)$ is perfect and hamiltonian. We also find the domination number of $\Gamma^*(\mathbb{Z}_n)$.

Theorem 3.1. Let $n = p_1^{\alpha_1}p_2^{\alpha_2} \cdots p_k^{\alpha_k}$, where p_i 's are distinct primes and $\alpha_i \geq 1$. Then $\Gamma^*(\mathbb{Z}_n)$ is Hamiltonian if and only if $k = 2$ and $\alpha_1 = \alpha_2$.

Proof. If $k = 2$ and $\alpha_1 = \alpha_2$, then $n = p_1^{\alpha_1} p_2^{\alpha_1}$. We now explicitly construct the hamiltonian circuit in $\Gamma^*(\mathbb{Z}_n)$:

$$\langle p_1 \rangle \sim \langle p_2 \rangle \sim \langle p_1^2 \rangle \sim \langle p_2^2 \rangle \sim \langle p_1^3 \rangle \sim \langle p_2^3 \rangle \sim \dots \sim \langle p_1^{\alpha_1} \rangle \sim \langle p_2^{\alpha_1} \rangle \sim \langle p_1 \rangle.$$

Conversely, let $\Gamma^*(\mathbb{Z}_n)$ be Hamiltonian. If possible, let $k \geq 3$. If $\alpha_i = 1$ for some i , then the graph has a vertex of degree 1 and hence it is not hamiltonian. Thus, we assume that $\alpha_i \geq 2$ for all i . Without loss of generality, let $\alpha_1 = \min\{\alpha_1, \alpha_2, \dots, \alpha_k\}$. Now, the vertices of the form $\langle p_2^{\alpha'_2} p_3^{\alpha'_3} \dots p_k^{\alpha'_k} \rangle$ are adjacent only to the vertices of the form $\langle p_1^{\alpha'_1} \rangle$, where $1 \leq \alpha'_i \leq \alpha_i$, i.e., we have $\alpha_2 \alpha_3 \dots \alpha_k$ distinct vertices of degree α_1 . As two vertices of the form $\langle p_2^{\alpha'_2} p_3^{\alpha'_3} \dots p_k^{\alpha'_k} \rangle$ are not adjacent, to complete a hamiltonian cycle, we need at least $\alpha_2 \alpha_3 \dots \alpha_k$ different vertices between the vertices of the form $\langle p_2^{\alpha'_2} p_3^{\alpha'_3} \dots p_k^{\alpha'_k} \rangle$. But, as $k \geq 3$, we have $\alpha_2 \alpha_3 \dots \alpha_k > \alpha_1$. This leads to a contradiction. Thus $k = 2$ and $n = p_1^{\alpha_1} p_2^{\alpha_2}$.

As earlier, we can assume that $\alpha_1, \alpha_2 \geq 2$. Let, if possible, $\alpha_1 \neq \alpha_2$. Without loss of generality, let $2 \leq \alpha_1 < \alpha_2$. Now, on any hamiltonian circuit in $\Gamma^*(\mathbb{Z}_n)$, between any two vertices of the form $\langle p_1^i \rangle$ and $\langle p_1^j \rangle$ we have a vertex of the form $\langle p_2^t \rangle$ and between any two vertices of the form $\langle p_2^i \rangle$ and $\langle p_2^j \rangle$ we have a vertex of the form $\langle p_1^t \rangle$. Thus any Hamiltonian circuit should consist of an alternating run of vertices of the form $\langle p_1^i \rangle$ and $\langle p_2^j \rangle$. However, as $\alpha_1 < \alpha_2$, we have more vertices of the form $\langle p_2^j \rangle$ than that of the form $\langle p_1^i \rangle$, a contradiction. Thus $\alpha_1 = \alpha_2$. \square

Theorem 3.2. *Let $n = p_1^{\alpha_1} p_2^{\alpha_2} \dots p_k^{\alpha_k}$, where p_i 's are distinct primes and $\alpha_i \geq 1$. Then $\Gamma^*(\mathbb{Z}_n)$ is perfect if and only if $k \leq 4$.*

Proof. We prove this by using Strong Perfect graph theorem. If $k \geq 5$, then there exists an induced 5-cycle in $\Gamma^*(\mathbb{Z}_n)$ as shown in Figure 1. Thus, in this case, $\Gamma^*(\mathbb{Z}_n)$ is not perfect. Let $k \leq 4$, i.e., n has at most 4 distinct prime factors p_1, p_2, p_3, p_4 . Let, if possible, $\Gamma^*(\mathbb{Z}_n)$ admits an induced odd cycle of length $t \geq 5$, say $\langle h_1 \rangle \sim \langle h_2 \rangle \sim \dots \sim \langle h_t \rangle \sim \langle h_1 \rangle$. From the non-adjacency relations, we get $\gcd(h_1, h_3), \gcd(h_1, h_4), \gcd(h_2, h_4), \gcd(h_2, h_5), \gcd(h_3, h_t) \neq 1$.

Let $p_1 \mid \gcd(h_1, h_3)$. Then $p_1 \mid h_1$ and $p_1 \mid h_3$. Again, as $\langle h_t \rangle \sim \langle h_1 \rangle$, we have $\gcd(h_1, h_t) = 1$, i.e., $p_1 \nmid h_t$.

Similarly, as $\langle h_3 \rangle \sim \langle h_4 \rangle$, we have $p_1 \nmid h_4$, i.e., $p_1 \nmid \gcd(h_1, h_4)$. Let $p_2 \mid \gcd(h_1, h_4)$. Then $p_2 \mid h_1$ and $p_2 \mid h_4$. Now as $\langle h_3 \rangle \sim \langle h_4 \rangle$, we conclude $p_2 \nmid h_3$.

Again, as $p_1, p_2 \mid h_1$ and $\langle h_1 \rangle \sim \langle h_2 \rangle$, $p_1, p_2 \nmid h_2$, i.e., $p_1, p_2 \nmid \gcd(h_2, h_4)$. Let $p_3 \mid \gcd(h_2, h_4)$. Then $p_3 \mid h_2$ and $p_3 \mid h_4$. As $\langle h_2 \rangle \sim \langle h_3 \rangle$, we have $p_3 \nmid h_3$.

Thus $p_1, p_2, p_3 \nmid \gcd(h_3, h_t)$. Let $p_4 \mid \gcd(h_3, h_t)$. Then $p_4 \mid h_3$ and $p_4 \mid h_t$. As $\langle h_2 \rangle \sim \langle h_3 \rangle$, $p_4 \nmid h_2$. Again, as $\langle h_4 \rangle \sim \langle h_5 \rangle$, we have $p_3 \nmid h_5$.

From the above situation, we get $p_1, p_2, p_3, p_4 \nmid \gcd(h_2, h_5)$. This is a contradiction, as $\gcd(h_2, h_5) \neq 1$ and $k \leq 4$. Thus $\Gamma^*(\mathbb{Z}_n)$ does not admit any induced odd cycle of length $t \geq 5$.

Let, if possible, the complement of $\Gamma^*(\mathbb{Z}_n)$ admits an induced odd cycle of length $t \geq 5$, say $\langle h_1 \rangle \sim \langle h_2 \rangle \sim \dots \sim \langle h_t \rangle \sim \langle h_1 \rangle$. Note that in the complement graph, two vertices $\langle h_i \rangle$ and $\langle h_j \rangle$ are non-adjacent/adjacent according as $\gcd(h_i, h_j)$ is equal/not equal to 1 respectively.

As $\langle h_1 \rangle \sim \langle h_2 \rangle$, we have $\gcd(h_1, h_2) \neq 1$. Let $p_1 \mid \gcd(h_1, h_2)$. Then $p_1 \mid h_1$ and $p_1 \mid h_2$. As $\gcd(h_1, h_3) = 1$, $p_1 \nmid h_3$, i.e., $p_1 \nmid \gcd(h_2, h_3)$. Similarly, we can conclude that p_1 does not divide any one of $\gcd(h_3, h_4), \gcd(h_4, h_5), \dots, \gcd(h_1, h_t)$.

Let $p_2 \mid \gcd(h_2, h_3)$. Then $p_2 \mid h_2$ and $p_2 \mid h_3$. As $\gcd(h_2, h_4) = 1$, we have $p_2 \nmid h_4$, i.e., p_2 does not divide $\gcd(h_3, h_4)$ and $\gcd(h_4, h_5)$. Similarly, as $\gcd(h_2, h_t) = 1$, we deduce $p_2 \nmid h_t$, i.e., $p_2 \nmid \gcd(h_1, h_t)$.

As $p_1, p_2 \nmid \gcd(h_3, h_4)$, let $p_3 \mid \gcd(h_3, h_4)$. Then $p_3 \mid h_3$ and $p_3 \mid h_4$. As $\gcd(h_1, h_3) = 1$, we have $p_3 \nmid h_1$, i.e., $p_3 \nmid \gcd(h_1, h_t)$. Similarly, as $\gcd(h_3, h_5) = 1$, we have $p_3 \nmid h_5$, i.e., $p_3 \nmid \gcd(h_4, h_5)$.

As $p_1, p_2, p_3 \nmid \gcd(h_4, h_5)$, let $p_4 \mid \gcd(h_4, h_5)$. Then $p_4 \mid h_4$ and $p_4 \mid h_5$. As $\gcd(h_1, h_4) = 1$, we have $p_4 \nmid h_1$, i.e., $p_4 \nmid \gcd(h_1, h_t)$.

Thus $p_1, p_2, p_3, p_4 \nmid \gcd(h_1, h_t)$. But this is a contradiction, as $\gcd(h_1, h_t) > 1$ and n has at most four distinct prime factors. Thus $\Gamma^*(\mathbb{Z}_n)^c$ does not admit an induced odd cycle of length $t \geq 5$.

Hence, by strong perfect graph theorem, the theorem follows. □




FIGURE 1. Induced 5-cycle in $\Gamma^*(\mathbb{Z}_n)$, for $k \geq 5$

Theorem 3.3. Let $n = p_1^{\alpha_1} p_2^{\alpha_2} \dots p_k^{\alpha_k}$, where p_i 's are distinct primes, $k \geq 2$ and $\alpha_i \geq 1$. Then the domination number of $\Gamma^*(\mathbb{Z}_n)$,

$$\gamma(\Gamma^*(\mathbb{Z}_n)) = \begin{cases} 1, & \text{if } n = p_1^{\alpha_1} p_2. \\ k, & \text{otherwise.} \end{cases}$$

Proof. Clearly $\{\langle p_1 \rangle, \langle p_2 \rangle, \dots, \langle p_k \rangle\}$ is a dominating set for $\Gamma^*(\mathbb{Z}_n)$ of size k . Thus $\gamma(\Gamma^*(\mathbb{Z}_n)) \leq k$.

Let $S = \{\langle x_1 \rangle, \langle x_2 \rangle, \dots, \langle x_{k-1} \rangle\}$ be a dominating set of $\Gamma^*(\mathbb{Z}_n)$ of size $k - 1$. Let $m = p_1 p_2 p_3 \dots p_k$. Out of the k vertices $\langle m/p_1 \rangle, \langle m/p_2 \rangle, \dots, \langle m/p_k \rangle$, at least one does not belong to S . Without loss of generality, let $\langle m/p_1 \rangle \notin S$ and $\langle m/p_1 \rangle \sim \langle x_1 \rangle$. Thus, by Lemma 2.1, $x_1 = p_1^{\alpha'_1}$, where $1 \leq \alpha'_1 \leq \alpha_1$. Thus $\langle x_1 \rangle$ is not adjacent to any of the $k - 1$ vertices $\langle m/p_2 \rangle, \langle m/p_3 \rangle, \dots, \langle m/p_k \rangle$. Again, by similar

argument, not all of these $k - 1$ vertices belong to S . Without loss of generality, let $\langle m/p_2 \rangle \notin S$ and $\langle m/p_2 \rangle \sim \langle x_2 \rangle$. Proceeding similarly, we get $x_2 = p_2^{\alpha'_2}$, where $1 \leq \alpha'_2 \leq \alpha_2$. Thus $\langle x_1 \rangle$ and $\langle x_2 \rangle$ are not adjacent to any of the $k - 2$ vertices $\langle m/p_3 \rangle, \dots, \langle m/p_k \rangle$. Continuing in this way, we get $x_i = p_i^{\alpha'_i}$ for $i = 1, 2, \dots, k - 1$. However, in that case, $\langle m/p_k \rangle$ neither belong to S nor adjacent to any element of S , a contradiction. Hence $\gamma(\Gamma^*(\mathbb{Z}_n)) = k$.

Note that the proof does not work if $k = 2$ and exactly one of the two powers is 1. Because in that case, one of $\langle m/p_1 \rangle$ and $\langle m/p_2 \rangle$ is not a vertex of $\Gamma^*(\mathbb{Z}_n)$, i.e., an isolated vertex of $\Gamma(\mathbb{Z}_n)$. If $k = 2$ and $n = p_1^{\alpha_1} p_2$, then $\langle p_2 \rangle$ dominates $\Gamma^*(\mathbb{Z}_n)$. \square

4. Isomorphisms

In this section, we discuss the conditions under which co-maximal subgroup graphs defined over different cyclic groups are isomorphic. For that, we start with the following definition.

Definition 4.1. *Two positive integers n and m are said to be of same prime-factorization type if $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$ and $m = q_1^{\beta_1} q_2^{\beta_2} \cdots q_k^{\beta_k}$ where p_i, q_i 's are primes and there exists $\sigma \in S_k$ such that $\alpha_i = \beta_{\sigma(i)}$ for $i = 1, 2, \dots, k$.*

Theorem 4.2. *Let n and m be two integers. Then $\Gamma(\mathbb{Z}_n) \cong \Gamma(\mathbb{Z}_m)$ if and only if m and n are of same prime-factorization type.*

Proof. If m and n are of same prime-factorization type, then the result is obvious. Let $\Gamma(\mathbb{Z}_n) \cong \Gamma(\mathbb{Z}_m)$, then as their clique numbers are equal, both m and n have same number of distinct prime factors. Let $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$ and $m = q_1^{\beta_1} q_2^{\beta_2} \cdots q_k^{\beta_k}$. Also as they have same number of isolated vertices, $\alpha_1 \cdot \alpha_2 \cdots \alpha_k = \beta_1 \cdot \beta_2 \cdots \beta_k$.

Without loss of generality, let $\alpha_1 = \min\{\alpha_1, \alpha_2, \dots, \alpha_k, \beta_1, \beta_2, \dots, \beta_k\}$. If possible, let $\alpha_1 \notin \{\beta_1, \beta_2, \dots, \beta_k\}$. Now, note that any vertex of the form $\langle p_2^{\alpha'_2} p_3^{\alpha'_3} \cdots p_k^{\alpha'_k} \rangle$ ($1 \leq \alpha'_i \leq \alpha_i$) in $\Gamma(\mathbb{Z}_n)$ is adjacent to only α_1 vertices, namely $\langle p_1 \rangle, \langle p_1^2 \rangle, \dots, \langle p_1^{\alpha_1} \rangle$. Thus $\Gamma(\mathbb{Z}_n)$ has $\alpha_2 \alpha_3 \cdots \alpha_k$ vertices of degree α_1 . As $\alpha_1 \leq \min\{\beta_1, \beta_2, \dots, \beta_k\}$ and $\alpha_1 \notin \{\beta_1, \beta_2, \dots, \beta_k\}$, from Theorem 2.2, it follows that $\Gamma(\mathbb{Z}_m)$ has no vertex of degree α_1 , a contradiction. Thus $\alpha_1 = \beta_i$ for some i . By suitable renaming, let $\alpha_1 = \beta_1$.

Again, without loss of generality, let $\alpha_2 = \min\{\alpha_2, \dots, \alpha_k, \beta_2, \dots, \beta_k\}$. If possible, let $\alpha_2 \notin \{\beta_2, \dots, \beta_k\}$. If $\alpha_2 \neq \beta_1$, then by a similar argument, $\Gamma(\mathbb{Z}_m)$ has no vertex of degree α_2 , a contradiction. Thus, we assume that $\alpha_2 = \beta_1 = \alpha_1$. Then $\Gamma(\mathbb{Z}_n)$ has $\alpha_2 \alpha_3 \cdots \alpha_k + \alpha_1 \alpha_3 \cdots \alpha_k$ of degree α_1 and $\Gamma(\mathbb{Z}_m)$ has $\beta_2 \beta_3 \cdots \beta_k$ of degree α_1 . As $\Gamma(\mathbb{Z}_n) \cong \Gamma(\mathbb{Z}_m)$, we have

$$\alpha_2 \alpha_3 \cdots \alpha_k + \alpha_1 \alpha_3 \cdots \alpha_k = \beta_2 \beta_3 \cdots \beta_k,$$

$$\text{i.e., } \alpha_3 \cdots \alpha_k (\alpha_1 + \alpha_2) = \frac{\alpha_1 \cdot \alpha_2 \cdots \alpha_k}{\beta_1} \quad (\text{as } \alpha_1 \cdot \alpha_2 \cdots \alpha_k = \beta_1 \cdot \beta_2 \cdots \beta_k)$$

$$\text{i.e., } \beta_1 (\alpha_1 + \alpha_2) = \alpha_1 \alpha_2, \text{ i.e., } 2\alpha_1^2 = \alpha_1^2, \text{ a contradiction.}$$

Thus $\alpha_2 = \beta_i$ for some $i \in \{2, 3, \dots, k\}$. By a suitable renaming, let $\alpha_2 = \beta_2$.

Proceeding this way, suppose in the $(l - 1)$ -th step, we get $\alpha_i = \beta_i$ for $i = 1, 2, \dots, l - 1$. Without loss of generality, let $\alpha_l = \min\{\alpha_l, \dots, \alpha_k, \beta_l, \dots, \beta_k\}$. If possible, let $\alpha_l \notin \{\beta_l, \dots, \beta_k\}$. If $\alpha_l \notin \{\beta_1, \beta_2, \dots, \beta_{l-1}\}$, then by a similar argument, $\Gamma(\mathbb{Z}_m)$ has no vertex of degree α_l , a contradiction. Thus, we assume that $\alpha_l \in \{\beta_1, \beta_2, \dots, \beta_{l-1}\}$. Let $\alpha_l = \beta_p = \beta_{p+1} = \dots = \beta_{l-1} = \alpha_p = \alpha_{p+1} = \dots = \alpha_{l-1}$ for some $1 \leq p \leq l - 1$.

Therefore, $\Gamma(\mathbb{Z}_n)$ has

$$\begin{aligned}
 & (\alpha_1 \alpha_2 \cdots \alpha_{p-1} \alpha_{p+1} \cdots \alpha_k) + (\alpha_1 \cdots \alpha_p \alpha_{p+1} \cdots \alpha_k) + \cdots + (\alpha_1 \cdots \alpha_{l-1} \alpha_{l+1} \cdots \alpha_k) \\
 &= \alpha_1 \cdots \alpha_k \left(\frac{1}{\alpha_p} + \frac{1}{\alpha_{p+1}} + \cdots + \frac{1}{\alpha_l} \right) \text{ vertices of degree } \alpha_l.
 \end{aligned}$$

Similarly, $\Gamma(\mathbb{Z}_m)$ has

$$\beta_1 \cdots \beta_k \left(\frac{1}{\beta_p} + \frac{1}{\beta_{p+1}} + \cdots + \frac{1}{\beta_{l-1}} \right) \text{ vertices of degree } \alpha_l.$$

Now, as $\Gamma(\mathbb{Z}_n) \cong \Gamma(\mathbb{Z}_m)$, we have

$$\alpha_1 \cdots \alpha_k \left(\frac{1}{\alpha_p} + \frac{1}{\alpha_{p+1}} + \cdots + \frac{1}{\alpha_l} \right) = \beta_1 \cdots \beta_k \left(\frac{1}{\beta_p} + \frac{1}{\beta_{p+1}} + \cdots + \frac{1}{\beta_{l-1}} \right)$$

$$\text{i.e., } \left(\frac{1}{\alpha_p} + \frac{1}{\alpha_{p+1}} + \cdots + \frac{1}{\alpha_l} \right) = \left(\frac{1}{\alpha_p} + \frac{1}{\alpha_{p+1}} + \cdots + \frac{1}{\alpha_{l-1}} \right) \Rightarrow \left(\frac{l - p + 1}{\alpha_l} \right) = \left(\frac{l - p}{\alpha_l} \right),$$

a contradiction. Thus, by a suitable renaming, we get $\alpha_l = \beta_l$, and hence by induction, the theorem follows. □

Acknowledgments

The first and third authors acknowledge the funding of DST-SERB-SRG Sanction no. *SRG/2019/000475* and *SRG/2019/000684*, Govt. of India. The second author is supported by the PhD fellowship of CSIR (File no. 08/155(0086)/2020-EMR-I), Govt. of India.

REFERENCES

- [1] S. Akbari, B. Miraftab and R. Nikandish, Co-maximal Graphs of Subgroups of Groups, *Canad. Math. Bull.*, **60** (2017) 12–25.
- [2] P. J. Cameron, Graphs defined on groups, *Int. J. Group Theory*, Article in Press. https://ijgt.ui.ac.ir/article_25608.html.
- [3] A. Das, M. Saha and S. Alkaseasbeh, On Co-Maximal Subgroup Graph of a Group, <https://arxiv.org/pdf/2103.14284.pdf>
- [4] B. Miraftab and R. Nikandish, Co-maximal Graphs of Two Generator Groups, *J. Algebra Appl.*, **18** (2019) pp. 13.

Manideepa Saha

Department of Mathematics, Presidency University, Kolkata, India

Email: manideepasaha1991@gmail.com

Sucharita Biswas

Department of Mathematics, Presidency University, Kolkata, India

Email: biswas.sucharita56@gmail.com

Angsuman Das

Department of Mathematics, Presidency University, Kolkata, India

Email: angsuman.maths@presiuniv.ac.in