



<http://ijgt.ui.ac.ir>



www.ui.ac.ir

A CHARACTERIZATION OF A_5 BY ITS AVERAGE ORDER

MARIUS TĂRNĂUCEANU

ABSTRACT. Let $o(G)$ be the average order of a finite group G . M. Herzog, P. Longobardi and M. Maj [M. Herzog, P. Longobardi and M. Maj, Another criterion for solvability of finite groups, *J. Algebra*, **597** (2022) 1-23.] showed that if G is non-solvable and $o(G) = o(A_5)$, then $G \cong A_5$. In this note, we prove that the equality $o(G) = o(A_5)$ does not hold for any finite solvable group G . Consequently, up to isomorphism, A_5 is determined by its average order.

1. Introduction

Given a finite group G , we denote by $\psi(G)$ the sum of element orders of G and by $o(G)$ the average order of G , that is

$$\psi(G) = \sum_{x \in G} o(x) \text{ and } o(G) = \frac{\psi(G)}{|G|}.$$

In the last years there has been a growing interest in studying the properties of these functions and their relations with the structure of G (see for example [1]-[8], [10],[11], [13] and [15]-[17]).

In [10], A. Jaikin-Zapirain uses the average order to determine a lower bound for the number of conjugacy classes of a finite p -group/nilpotent group. He also suggests the following question: “Let G be a finite (p -)group and N be a normal (abelian) subgroup of G . Is it true that $o(G) \geq o(N)^{\frac{1}{2}}$?”. Recently, E.I. Khukhro, A. Moretó and M. Zarrin proved the following result (see [11, Theorem 1.2]):

Theorem A. *Let $c > 0$ be a real number and $p \geq \frac{3}{c}$ be a prime. Then there exists a finite p -group G with a normal abelian subgroup N such that $o(G) < o(N)^c$.*

Keywords: average order, sum of element orders, solvable group.

MSC(2010): Primary: 20D60; Secondary: 20D10, 20F16.

Communicated by Alireza Abdollahi.

Received: 6 August 2023, Accepted: 30 December 2023.

Cite this article: M. Tărnăuceanu, A characterization of A_5 by its average order, Int. J. Group Theory, x no. x (202x) xx-xx.

<http://dx.doi.org/10.22108/ijgt.2023.138637.1866> .

Note that Theorem A provides a negative answer to Jaikin-Zapirain's question even if we replace the exponent $\frac{1}{2}$ with any positive real number c . In the same paper [11], the authors posed the following conjecture:

Conjecture. *Let G be a finite group and suppose that $o(G) < \frac{211}{60} = o(A_5)$. Then G is solvable.*

It has been confirmed by M. Herzog, P. Longobardi and M. Maj [7].

Theorem B. *Let G be a finite group. If $o(G) < \frac{211}{60}$, then G is solvable. Moreover, if G is a non-solvable finite group, we have $o(G) = \frac{211}{60}$ if and only if $G \cong A_5$.*

This constitutes the starting point for the current work. Our main result is as follows.

Theorem 1.1. *If G is a finite solvable group, then $o(G) \neq \frac{211}{60}$.*

It is now clear that Theorems B and 1.1 lead to the following characterization of A_5 .

Corollary 1.2. *Up to isomorphism, A_5 is the unique finite group G such that $o(G) = \frac{211}{60}$.*

Note that the same result was obtained by M. Herzog, P. Longobardi and M. Maj [8]. Our proof is different from that in [8] and based upon the next theorem.

Theorem C. *Let G be a finite group and $n_d(G)$ be the number of elements of order d in G , $\forall d \in \mathbb{N}$. Then the following statements hold:*

1) [12] *If p is a prime divisor of $|G|$ and G is not a p -group, then*

$$n_p(G) \leq \frac{p}{p+1} |G| - 1.$$

2) [14] *If G is non-abelian, then*

$$n_2(G) \leq \frac{3}{4} |G| - 1.$$

Most of our notation is standard and will usually not be repeated here. Elementary notions and results on groups can be found in [9].

2. Proofs of the main results

We start with the following easy but important lemma.

Lemma 2.1. *Let G be a finite group of order n and $n_d(G)$ be the number of elements of order d in G , $\forall d \in \mathbb{N}^*$. If $o(G) = \frac{211}{60}$, then*

$$(2.1) \quad 4n_2(G) + 3n_3(G) + 2n_4(G) + n_5(G) \geq \frac{149}{60} n - 5.$$

Proof. Since $n = \sum_{d \in \mathbb{N}^*} n_d(G)$, we deduce that

$$\begin{aligned} \psi(G) &= 1 + 2n_2(G) + \dots + 6n_6(G) + \dots \\ &\geq 1 + 2n_2(G) + \dots + 5n_5(G) + 6(n_6(G) + n_7(G) + \dots) \\ &= 1 + 2n_2(G) + \dots + 5n_5(G) + 6(n - 1 - n_2(G) - \dots - n_5(G)) \\ &= 6n - 5 - (4n_2(G) + 3n_3(G) + 2n_4(G) + n_5(G)). \end{aligned}$$

Since $o(G) = \frac{211}{60}$, it follows that $\psi(G) = \frac{211}{60}n$. Therefore we have

$$\frac{211}{60}n \geq 6n - 5 - (4n_2(G) + 3n_3(G) + 2n_4(G) + n_5(G)),$$

which is equivalent to (1), completing the proof. □

Our second lemma shows that the average order of a finite nilpotent group which is not a p -group cannot be certain rational numbers.

Lemma 2.2. *Let G be a finite nilpotent group. If G is not a p -group, then $o(G) \notin \left\{ \frac{61}{15}, \frac{211}{60} \right\}$.*

Proof. We write G as a direct product of its Sylow subgroups

$$G = G_1 \times \dots \times G_k,$$

where $k \geq 2$ and $G_i \in \text{Syl}_{p_i}(G)$, for all $i = 1, \dots, k$. Since the function o is multiplicative, we get

$$o(G) = o(G_1) \dots o(G_k).$$

Let $|G_i| = p_i^{n_i}$, $i = 1, \dots, k$, and assume that $p_1 < \dots < p_k$. Then for each i we have

$$o(G_i) \geq \frac{1 + p_i(p_i^{n_i} - 1)}{p_i^{n_i}} = p_i - \frac{p_i - 1}{p_i^{n_i}} \geq p_i - \frac{p_i - 1}{p_i} > 1.$$

For $k \geq 3$, we get

$$o(G) \geq o(G_1)o(G_2)o(G_3) \geq \left(2 - \frac{1}{2}\right)\left(3 - \frac{2}{3}\right)\left(5 - \frac{4}{5}\right) = \frac{147}{10} > \frac{61}{15} > \frac{211}{60}.$$

If $k = 2$ and $(p_1, p_2) \neq (2, 3)$, then

$$o(G) = o(G_1)o(G_2) \geq \min\left\{\left(3 - \frac{2}{3}\right)\left(5 - \frac{4}{5}\right), \left(2 - \frac{1}{2}\right)\left(5 - \frac{4}{5}\right)\right\} = \frac{63}{10} > \frac{61}{15}.$$

Thus, we can suppose that $(p_1, p_2) = (2, 3)$. If $(n_1, n_2) \neq (1, 1)$, then

$$o(G) = o(G_1)o(G_2) \geq \min\left\{\left(2 - \frac{1}{4}\right)\left(3 - \frac{2}{3}\right), \left(2 - \frac{1}{2}\right)\left(3 - \frac{2}{9}\right)\right\} = \frac{49}{12} > \frac{61}{15}.$$

Finally, for $(n_1, n_2) = (1, 1)$ we have $G \cong C_6$ and so

$$o(G) = \frac{7}{2} \notin \left\{ \frac{61}{15}, \frac{211}{60} \right\},$$

as desired. □

We are now able to prove our main result. Note that a search with GAP shows that there is no solvable group G of order less than 1000 such that $o(G) = \frac{211}{60}$.

Proof of Theorem 1.1. Assume that there exists a finite solvable group G of order n with $o(G) = \frac{211}{60}$. Then $60\psi(G) = 211n$, implying that $n = 2^{2\alpha_2}3^{\alpha_2}5^{\alpha_3} \dots$ with $\alpha_2, \alpha_3 \geq 1$. Let H be a Sylow 2-subgroup of G . We distinguish the following two cases.

Case 1. $H \cong C_4$ Then G has a normal 2-complement K of order $m = 3^{\alpha_2}5^{\alpha_3} \dots$. First of all, we prove that

$$(2.2) \quad n_4(G) = 2m = \frac{1}{2}n.$$

Suppose that $n_4(G) < 2m$. Since $\frac{n_4(G)}{2}$ divides m , it follows that $\frac{n_4(G)}{2} \leq \frac{m}{3}$ and so $n_4(G) \leq \frac{2}{3}m = \frac{1}{6}n$. This implies that $n_2(G) \leq \frac{1}{12}n$. Also, since all elements of odd order of G are contained in K , we get $n_3(G), n_5(G) < m = \frac{1}{4}n$. Thus we have

$$4n_2(G) + 3n_3(G) + 2n_4(G) + n_5(G) < \left(\frac{1}{3} + \frac{3}{4} + \frac{1}{3} + \frac{1}{4}\right)n = \frac{5}{3}n,$$

which together with (1) lead to

$$\frac{149}{60}n - 5 < \frac{5}{3}n,$$

that is $n \leq 6$, a contradiction.

Next we prove that

$$(2.3) \quad n_2(G) = m = \frac{1}{4}n.$$

Suppose that $n_2(G) < m$. Since every subgroup of order 2 of G is contained in at least three Sylow 2-subgroups, we infer that $n_2(G) \leq \frac{m}{3} = \frac{1}{12}n$. Similarly, we get

$$4n_2(G) + 3n_3(G) + 2n_4(G) + n_5(G) < \left(\frac{1}{3} + \frac{3}{4} + 1 + \frac{1}{4}\right)n = \frac{7}{3}n$$

and so

$$\frac{149}{60}n - 5 < \frac{7}{3}n$$

by (1). Thus $n \leq 33$, a contradiction.

The equalities (2) and (3) show that any two Sylow 2-subgroups of G intersect trivially and any element of G of even order is a 2-element. We deduce that G is a Frobenius group and consequently its kernel K is nilpotent. On the other hand, we have

$$\frac{211}{60}n = \psi(G) = \psi(K) + 2n_2(G) + 4n_4(G) = \psi(K) + \frac{5}{2}n,$$

implying that

$$\psi(K) = \frac{61}{60}n = \frac{61}{15}m,$$

or equivalently

$$o(K) = \frac{61}{15}.$$

This contradicts Lemma 2.2.

Case 2. $H \cong C_2^2$ Since G is solvable, it has a Hall 2-subgroup K of order $m = 3^{\alpha_2}5^{\alpha_3} \dots$. We have the next two subcases.

Subcase 2.1. K is normal in G

Then K contains all elements of odd order of G . By using Theorem C, 1), for $p = 3$, it follows that $n_3(G) \leq \frac{3}{4}m = \frac{3}{16}n$. Also, we have $n_3(G) + n_5(G) \leq m - 1 = \frac{n}{4} - 1$. Then

$$3n_3(G) + n_5(G) = 2n_3(G) + (n_3(G) + n_5(G)) \leq \frac{3}{8}n + \frac{1}{4}n - 1 = \frac{5}{8}n - 1,$$

implying that

$$4n_2(G) + 3n_3(G) + 2n_4(G) + n_5(G) \leq 4n_2(G) + \frac{5}{8}n - 1.$$

From (1) we get

$$(2.4) \quad n_2(G) \geq \frac{223}{480}n - 1.$$

Let $H_i, i = 1, 2, 3$, be the subgroups of order 2 of H . Then $G = KH_1 \cup KH_2 \cup KH_3$ and so $n_2(G) = 3n_2(KH_i)$, for any $i = 1, 2, 3$. On the other hand, we have $n_2(KH_i) \mid m$ by Sylow's theorems. If $n_2(KH_i) < m$, then $n_2(KH_i) \leq \frac{m}{3}$, which lead to

$$(2.5) \quad n_2(G) \leq m = \frac{1}{4}n.$$

Now, from (4) and (5), we obtain

$$\frac{223}{480}n - 1 \leq \frac{1}{4}n,$$

that is $n \leq 4$, a contradiction. Thus $n_2(KH_i) = m$, for all $i = 1, 2, 3$, and $n_2(G) = 3m = \frac{3}{4}n$. Then, by Theorem C, 2), it follows that G is abelian. Since $o(G) = \frac{211}{60}$, this contradicts Lemma 2.2.

Subcase 2.2. K is not normal in G

Then K has two or four conjugates in G . Let $C = \text{Core}_G(K)$. Since $[G : C]$ divides $4! = 24$, we get $[G : C] = 12$ and so $[K : C] = 3$. We observe that A_4 is the unique group of order 12 containing more than one subgroup of order 3. Thus $G/C \cong A_4$ and K has four conjugates in G , say $K_i, i = 1, \dots, 4$. Then G possesses

$$\left| \bigcup_{i=1}^4 K_i \right| = m + 3 \left(m - \frac{m}{3} \right) = 3m$$

elements of odd order. Also, we have

$$(2.6) \quad \begin{aligned} n_3(G) &= n_3 \left(\bigcup_{i=1}^4 K_i \right) \leq n_3(K_1) + 3 \left| \bigcup_{i=2}^4 (K_i \setminus K_1) \right| \\ &\leq \frac{3m}{4} + 3 \left(m - \frac{m}{3} \right) = \frac{11}{4}m = \frac{11}{16}n \end{aligned}$$

by applying Theorem C, 1), for the group K_1 and $p = 3$.

Assume that $n_2(G) = m$. Then the number s_2 of Sylow 2-subgroups of G is at least $\frac{m}{3}$. Since all Sylow 2-subgroups of G are contained in CH and $|CH| = 4 \cdot \frac{m}{3}$, we get $s_2 \mid \frac{m}{3}$ and therefore $s_2 = \frac{m}{3}$. This implies again that G is a Frobenius group, a contradiction. Thus

$$n_2(G) \leq \frac{m}{3} = \frac{n}{12}.$$

Since C includes all Sylow 5-subgroups of G , it results that

$$n_5(G) \leq |C| - 1 = \frac{m}{3} - 1 = \frac{n}{12} - 1.$$

Then

$$4n_2(G) + 3n_3(G) + 2n_4(G) + n_5(G) \leq \frac{n}{3} + 3n_3(G) + \frac{n}{12} - 1 = 3n_3(G) + \frac{5}{12}n - 1$$

which together with (1) lead to

$$3n_3(G) \geq \left(\frac{149}{60} - \frac{5}{12} \right) n - 4,$$

that is

$$(2.7) \quad n_3(G) \geq \frac{31}{45}n - \frac{4}{3}.$$

Now, from (6) and (7), we obtain

$$\frac{31}{45}n - \frac{4}{3} \leq \frac{11}{16}n,$$

that is $n \leq 960$, a contradiction. □

Acknowledgments

The author is grateful to the reviewer for remarks which improve the previous version of the paper.

REFERENCES

- [1] H. Amiri and S. M. Jafarian Amiri, Sum of element orders on finite groups of the same order, *J. Algebra Appl.*, **10** no. 2 (2011) 187–190.
- [2] H. Amiri, S. M. Jafarian Amiri and I. M. Isaacs, Sums of element orders in finite groups, *Comm. Algebra*, **37** no. 9 (2009) 2978–2980.
- [3] M. Baniasad Azad and B. Khosravi, A criterion for solvability of a finite group by the sum of element orders, *J. Algebra*, **516** (2018) 115–124.
- [4] M. Baniasad Azad and B. Khosravi, On two conjectures about the sum of element orders, *Canad. Math. Bull.*, **65** no. 1 (2022) 30–38.
- [5] M. Herzog, P. Longobardi and M. Maj, Two new criteria for solvability of finite groups, *J. Algebra*, **511** (2018) 215–226.
- [6] M. Herzog, P. Longobardi and M. Maj, An exact upper bound for sums of element orders in non-cyclic finite groups, *J. Pure Appl. Algebra*, **222** no. 7 (2018) 1628–1642.
- [7] M. Herzog, P. Longobardi and M. Maj, Another criterion for solvability of finite groups, *J. Algebra*, **597** (2022) 1–23.

- [8] M. Herzog, P. Longobardi and M. Maj, On groups with average element orders equal to the average order of the alternating group of degree 5, to appear in *Glas. Mat. III. Ser.*, (2023).
- [9] I. M. Isaacs, *Finite group theory*, Graduate Studies in Mathematics, **92**, American Mathematical Society, Providence, RI, 2008
- [10] A. Jaikin-Zapirain, On the number of conjugacy classes of finite nilpotent groups, *Adv. Math.*, no. 3 **227** (2011) 1129–1143.
- [11] E. I. Khukhro, A. Moretó and M. Zarrin, The average element order and the number of conjugacy classes of finite groups, *J. Algebra*, **569** (2021) 1–11.
- [12] T. J. Laffey, The number of solution of $x^p = 1$ in a finite group, *Math. Proc. Cambridge Philos. Soc.*, **80** no. 2 (1976) 229–231.
- [13] M. S. Lazorec and M. Tărnăuceanu, On the average order of a finite group, *J. Pure Appl. Algebra*, **227** no. 4 (2023) 9 pp.
- [14] G. A. Miller, On the minimum number of operators whose orders exceed two in any finite group, *Bull. Amer. Math. Soc.*, **13** no. 5 (1907) 235–239.
- [15] M. Tărnăuceanu, Detecting structural properties of finite groups by the sum of element orders, *Israel J. Math.*, **238** no. 2 (2020) 629–637.
- [16] M. Tărnăuceanu, A criterion for nilpotency of a finite group by the sum of element orders, *Comm. Algebra*, **49** no. 4 (2021) 1571–1577.
- [17] M. Tărnăuceanu, Another criterion for supersolvability of finite groups, *J. Algebra*, **604** (2022) 682–693.
- [18] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.11.0*, 2020, <https://www.gap-system.org>.

Marius Tărnăuceanu

Faculty of Mathematics, Al.I. Cuza University, Iași, Romania

Email: tarnauc@uaic.ro