



<https://toc.ui.ac.ir>

---

**Transactions on Combinatorics**

ISSN (print): 2251-8657, ISSN (on-line): 2251-8665

Vol. x No. x (201x), pp. xx-xx.

© 20xx University of Isfahan

---



[www.ui.ac.ir](http://www.ui.ac.ir)

## ON THE INDICES OF CERTAIN GRAPH PRODUCTS

ISHITA SARKAR AND MANJUNATH NANJAPPA\*<sup>ORCID</sup>

**ABSTRACT.** Molecular descriptors are numerical graph invariants that are used to study the chemical structure of molecules. In this paper, we determine the upper bound of the Sombor index based on four operations involving the subdivision graph, semi-total point graph, semi-total line graph, and total graph related to the lexicographic and tensor product. The exact expressions of the first reformulated Zagreb index and the second hyper-Zagreb index of the tensor product are formulated on the basis of the four significant graphs. Further, the descriptors for certain standard graphs are obtained and the graphical comparison for the first reformulated Zagreb index has also been illustrated to understand the result better.

### 1. Introduction

Topology of a molecule is primarily a non-mathematical that reflects various features of a molecular network. Numerous quantitative molecular attributes are frequently represented by precise numerals. To analyze topological invariants, a chemical compound must be amended to a molecular graph, with the atoms in the molecule corresponding to vertices and the atomic connections represented as edges. The topological indices are beneficial in many fields, but degree-based invariants are notably essential in the arena of chemical graph theory.

$V(H)$  and  $E(H)$  denotes the vertex and edge set for a molecular graph  $H = (V, E)$ , with their cardinality as  $p$  and  $q$  respectively. The degree of vertex  $z$  of  $H$  is denoted by  $d(z)$  and  $e = yz$  is a

---

Keywords: Sombor index, Lexicographic product, Tensor product.

MSC(2010): Primary: 05C09; Secondary: 05C76, 05C92.

Communicated by Alireza Abdollahi.

Article Type: Research Paper.

\*Corresponding author.

Received: 22 December 2022, Accepted: 16 July 2024, Published Online: 17 September 2024.

**Cite this article:** I. Sarkar and M. Nanjappa, On the indices of certain graph products, *Trans. Comb.*, **xx** no. x (20xx) xx-xx.

<http://dx.doi.org/10.22108/toc.2024.136176.2034> .

representation of the edge linking the nodes  $y$  and  $z$ . The line graph of  $H$ ,  $L(H)$  is the graph wherein the edges of  $H$  correspond to vertices of  $L(H)$  and two edges of  $L(H)$  are adjacent if and only if they are incident in  $H$ .  $\delta_H$  and  $\Delta_H$  indicates minimum and maximum degree for the graph,  $H$  respectively.

Initiated by Gutman [18, 17], the First ( $M_1$ ) and the Second ( $M_2$ ) Zagreb Index are the widely known degree based topological descriptors for the specification of  $\pi$ -electron energy of the molecules. The study on first general Zagreb index,  $M_\alpha$  ( $\alpha \in \mathbb{R}$ ,  $\alpha \neq 0, 1$ ) was initiated by the implementation of particular generalizations by Xueliang Li and Jie Zheng [25].

$$M_\alpha(H) = \sum_{z \in V(H)} (d_H(z))^\alpha$$

The sum-connectivity index,  $\chi$  was proposed by Bo Zhou and Nenad Trinajstić. Following that, several properties and discrete constraints for the associated invariant were determined in [40].

$$\chi(H) = \sum_{yz \in E(H)} (d_H(y) + d_H(z))^{-1/2}$$

On generalizing sum-connectivity index with the first Zagreb index laid the foundation of research on another invariant called the general sum-connectivity index,  $\chi_\alpha$  ( $\alpha \in \mathbb{R}$ ) [41]. Significant core concepts for the descriptor have also been investigated in regard to specific graph operations [1].

$$\chi_\alpha(H) = \sum_{yz \in E(H)} (d_H(y) + d_H(z))^\alpha$$

Detailed study on another descriptor, the forgotten topological index have also been acted upon in papers [12]. Various graph operational expressions have been determined with reference to the index [8].

$$\mathbf{F}(H) = \sum_{z \in V(H)} d_H(z)^3 \text{ or } \sum_{yz \in E(H)} (d_H(y)^2 + d_H(z)^2)$$

The study on first hyper Zagreb index ( $HM_1$ ) was initiated in [34] and specific outcomes with reference to the mathematical graph operational procedures were obtained. In addition, the formulation of second hyper Zagreb index ( $HM_2$ ) was proposed in [11] and computational techniques pertaining to various classes of graphs have been established [13].

$$HM_1(H) = \sum_{yz \in E(H)} (d_H(y) + d_H(z))^2 \text{ and } HM_2(H) = \sum_{yz \in E(H)} (d_H(y) \cdot d_H(z))^2$$

In relation to the edge degrees, reformulations of the Zagreb indices have been identified in [29] as

$$EM_1(H) = \sum_{e \in E(H)} d_H(e)^2 \text{ and } EM_2(H) = \sum_{e \sim f \in E(H)} d_H(e) d_H(f)$$

Lately, the theoretical implications of Sombor index( $SO$ ) was postulated by Gutman[14] and further work on the graph index has been acted upon[6, 15, 21, 37].

$$SO(H) = \sum_{yz \in E(H)} \sqrt{d_H(y)^2 + d_H(z)^2}$$

The analysis of Sombor index related to some graph operations have been resolved in [2, 23]. Also recently, reformulation of the Sombor index have been determined in terms of edge degrees in [19].

On a connected graph  $H$ , [10, 39, 32] defines the graphs related as:

- (i)  $S(H)$  or subdivision graph is generated by replacing every edge of the graph with a vertex of degree two, keeping the original vertices unchanged.
- (ii)  $R(H)$  or semi total point graph is the graph including the edges of  $S(H)$  along with the edges of  $H$ .
- (iii)  $Q(H)$  or semi total line graph is the one including edges of  $S(H)$  along with the edges of  $L(H)$ .
- (iv)  $T(H)$  or total graph includes edges of  $S(H)$ ,  $L(H)$  along with the edges of  $H$ .




FIGURE 1. Base graph and some of its derived graphs

The lexicographic product of graphs  $H_1$  and  $H_2$ , denoted as  $H_1[H_2]$ [38], is a graph wherein  $V(H_1[H_2]) = V(H_1) \times V(H_2)$  and  $(y_1, z_1)$  is adjacent to  $(y_2, z_2)$  of  $H_1[H_2]$  if and only if either  $y_1 = y_2$  and  $z_1$  is adjacent to  $z_2$  or  $y_1$  is adjacent to  $y_2$ . The tensor product  $H_1 \otimes H_2$ [22] for the graphs  $H_1$  and  $H_2$  possesses the vertex set as  $V(H_1) \times V(H_2)$  wherein  $(y_1, z_1)$  is incident on  $(y_2, z_2)$  whenever in  $H_1$ ,  $y_1$  is adjacent to  $y_2$  and in  $H_2$ ,  $z_1$  is adjacent to  $z_2$ . Works pertaining to the structure-invariants of the lexicographic and tensor products have been carried out[3, 36, 7, 31, 9, 5]. Works on various graph join and corona product variants have been established for  $SK$  indices in [28]. Also, findings related to other significant graph invariants of graph operations have also been investigated [27, 35, 20, 24, 16, 26, 30].

## 2. Methodology

The notion of the graph operational products is to obtain many pairs of co-spectral graphs. The operational procedure begins with simple graph and results in complex network structures from the base graphs. The  $F$ -lexicographic product of  $H_1$  and  $H_2$  for  $F \in \{S, R, Q, T\}$  denoted by  $H_1[H_2]_F$  [33] is a graph with  $V(H_1[H_2]_F) = (V(H_1) \cup E(H_1)) \times V(H_2)$  and  $u = (y_1, z_1)$  and  $v = (y_2, z_2)$  of  $H_1[H_2]$  are adjacent if and only if either  $[y_1 y_2 \in E(F(H_1)) \text{ and } z_1, z_2 \in V(H_2)]$  or  $[y_1 = y_2 \in V(H_1) \text{ and } z_1 z_2 \in E(H_2)]$ . Below we have representations of  $P_4[P_2]_S$ ,  $P_4[P_2]_R$ ,  $P_4[P_2]_Q$  and  $P_4[P_2]_T$  in the figure (2).




FIGURE 2.  $P_4[P_2]_F$

Assume  $F$  belongs to one among the symbols  $S$ ,  $R$ ,  $Q$  or  $T$ . The  $F$ -tensor product  $H_1 \otimes_F H_2$  [4] depicts a graph having  $V(H_1 \otimes_F H_2) = (V(H_1) \cup E(H_1)) \times V(H_2)$  and vertices  $(y_1, z_1)$  and  $(y_2, z_2)$  of  $H_1 \otimes_F H_2$  are adjacent if and only if  $y_1$  is incident on  $y_2$  in  $E(F(H_1))$  and  $z_1$  is incident on  $z_2$  in  $H_2$ . Below we have representations of  $P_4 \otimes_S P_2$ ,  $P_4 \otimes_R P_2$ ,  $P_4 \otimes_Q P_2$  and  $P_4 \otimes_T P_2$  in the figure (3).




FIGURE 3.  $P_4 \otimes_F P_2$

**2.1. Sombor index based on four operations related to the lexicographic and tensor product.**

In this section, we obtain the upper bounds of Sombor index based on the four operations related to the lexicographic and tensor product of two graphs. In addition, we have carried out the determination of the upper bound expressions for the descriptor of some standard graphs like path and cycle graphs.

**Theorem 2.1.** *Assume  $H_1(p_1, q_1)$  and  $H_2(p_2, q_2)$  be two arbitrary graphs. Then,*

$$SO(H_1[H_2]_S) \leq p_2^3(M_1(H_1) + 4q_1) + p_1SO(H_2) + (2p_2)^{1/2}(p_2M_{3/2}(H_1)M_{1/2}(H_2) + M_{1/2}(H_1)\chi_{1/2}(H_2)) + 2^{3/2}(2^{1/2} + 1)p_2q_1q_2$$

*Equality holds if and only if the components of the graph are isolated vertices.*

*Proof.*

$$\begin{aligned} SO(H_1[H_2]_S) &= \sum_{(y_1, z_1)(y_2, z_2) \in E(H_1[H_2]_S)} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\ &= \sum_{y_1=y_2 \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\ &+ \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{y_1 y_2 \in E(S(H_1))} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\ &= S_1 + S_2 \end{aligned}$$

$$\begin{aligned}
S_1 &= \sum_{y_1=y_2 \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\
&= \sum_{y \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{(p_2 d_{H_1}(y) + d_{H_2}(z_1))^2 + (p_2 d_{H_1}(y) + d_{H_2}(z_2))^2} \\
&= \sum_{y \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{p_2^2 d_{H_1}(y)^2 + d_{H_2}(z_1)^2 + 2p_2 d_{H_1}(y) d_{H_2}(z_1) + p_2^2 d_{H_1}(y)^2 + d_{H_2}(z_2)^2 + 2p_2 d_{H_1}(y) d_{H_2}(z_2)} \\
&\leq p_1 SO(H_2) + (2p_2)^{1/2} M_{1/2}(H_1) \chi_{1/2}(H_2) + 2^{3/2} p_2 q_1 q_2 \\
S_2 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{y_1 y_2 \in E(S(H_1))} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\
&= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y \in V(H_1) \\ a \in V(S(H_1)) \setminus V(H_1) \\ y \sim a}} \sqrt{d(y, z_1)^2 + d(a, z_2)^2} \\
&= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y \in V(H_1) \\ a \in V(S(H_1)) \setminus V(H_1) \\ y \sim a}} \sqrt{(p_2 d_{H_1}(y) + d_{H_2}(z_1))^2 + (2p_2)^2} \\
&\leq p_2^3 M_1(H_1) + (2p_2^3)^{1/2} M_{3/2}(H_1) M_{1/2}(H_2) + 4p_2 q_1 q_2 + 4p_2^3 q_1
\end{aligned}$$

From all the computations,

$$\begin{aligned}
SO(H_1[H_2]_S) &\leq p_2^3 (M_1(H_1) + 4q_1) + p_1 SO(H_2) + (2p_2)^{1/2} (p_2 M_{3/2}(H_1) M_{1/2}(H_2) + M_{1/2}(H_1) \chi_{1/2}(H_2)) \\
&\quad + 2^{3/2} (2^{1/2} + 1) p_2 q_1 q_2
\end{aligned}$$

Equality holds if and only if the components of the graph are isolated vertices. This concludes the result.  $\square$

**Theorem 2.2.** Assume  $H_1(p_1, q_1)$  and  $H_2(p_2, q_2)$  be two arbitrary graphs. Then,

$$\begin{aligned}
SO(H_1[H_2]_R) &\leq 2p_2^3 (SO(H_1) + M_1(H_1)) + p_1 SO(H_2) + 2p_2^{1/2} \left[ M_{1/2}(H_1) \chi_{1/2}(H_2) + p_2 M_{3/2}(H_1) \right. \\
&\quad \left. M_{1/2}(H_2) \right] + 4(2)^{1/2} (2^{1/2} + 1) p_2 q_1 q_2 + 4p_2^3 q_1 + (8p_2^5 \Delta_{H_1} \Delta_{H_2})^{1/2} q_1
\end{aligned}$$

Equality holds if and only if the components of the graph are isolated vertices.

*Proof.*

$$\begin{aligned}
SO(H_1[H_2]_R) &= \sum_{(y_1, z_1)(y_2, z_2) \in E(H_1[H_2]_R)} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\
&= \sum_{y_1=y_2 \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\
&\quad + \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{y_1 y_2 \in E(R(H_1))} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\
&= R_1 + R_2
\end{aligned}$$

$$\begin{aligned}
 R_1 &= \sum_{y_1=y_2 \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\
 &= \sum_{y_1=y_2 \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{(p_2 d_{R(H_1)}(y_1) + d_{H_2}(z_1))^2 + (p_2 d_{R(H_1)}(y_1) + d_{H_2}(z_2))^2} \\
 &= \sum_{y \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{(2p_2 d_{H_1}(y) + d_{H_2}(z_1))^2 + (2p_2 d_{H_1}(y) + d_{H_2}(z_2))^2} \\
 &\leq p_1 SO(H_2) + 2(p_2)^{1/2} M_{1/2}(H_1) \chi_{1/2}(H_2) + 4(2)^{1/2} p_2 q_1 q_2
 \end{aligned}$$

$$\begin{aligned}
 R_2 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{y_1 y_2 \in E(R(H_1))} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\
 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{y_1 y_2 \in E(H_1)} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\
 &\quad + \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(R(H_1)) \\ y_1 \in V(H_1) \\ y_2 \in V(R(H_1)) \setminus V(H_1)}} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\
 &= R'_2 + R''_2
 \end{aligned}$$

$$\begin{aligned}
 R'_2 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{y_1 y_2 \in E(H_1)} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\
 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{y_1 y_2 \in E(H_1)} \sqrt{(p_2 d_{R(H_1)}(y_1) + d_{H_2}(z_1))^2 + (p_2 d_{R(H_1)}(y_2) + d_{H_2}(z_2))^2} \\
 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{y_1 y_2 \in E(H_1)} \sqrt{(2p_2 d_{H_1}(y_1) + d_{H_2}(z_1))^2 + (2p_2 d_{H_1}(y_2) + d_{H_2}(z_2))^2} \\
 &\leq 2p_2^3 SO(H_1) + (8p_2^5 \Delta_{H_1} \Delta_{H_2})^{1/2} q_1 + 4p_2 q_1 q_2
 \end{aligned}$$

$$\begin{aligned}
 R''_2 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(R(H_1)) \\ y_1 \in V(H_1) \\ y_2 \in V(R(H_1)) \setminus V(H_1)}} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\
 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(R(H_1)) \\ y_1 \in V(H_1) \\ y_2 \in V(R(H_1)) \setminus V(H_1)}} \sqrt{(p_2 d_{R(H_1)}(y_1) + d_{H_2}(z_1))^2 + (p_2 d_{R(H_1)}(y_2))^2} \\
 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(R(H_1)) \\ y_1 \in V(H_1) \\ y_2 \in V(R(H_1)) \setminus V(H_1)}} \sqrt{(2p_2 d_{H_1}(y_1) + d_{H_2}(z_1))^2 + (2p_2)^2} \\
 &\leq 2p_2^3 M_1(H_1) + 2p_2^{3/2} M_{3/2}(H_1) M_{1/2}(H_2) + 4p_2 q_1 q_2 + 4p_2^3 q_1
 \end{aligned}$$

Hence from all the computations,

$$SO(H_1[H_2]_R) \leq 2p_2^3(SO(H_1) + M_1(H_1)) + p_1SO(H_2) + 2p_2^{1/2} \left[ M_{1/2}(H_1)\chi_{1/2}(H_2) + p_2M_{3/2}(H_1)M_{1/2}(H_2) \right] + 4(2)^{1/2}(2^{1/2} + 1)p_2q_1q_2 + 4p_2^3q_1 + (8p_2^5\Delta_{H_1}\Delta_{H_2})^{1/2}q_1$$

Equality holds if and only if the components of the graph are isolated vertices. This concludes the result.  $\square$

**Theorem 2.3.** Assume  $H_1(p_1, q_1)$  and  $H_2(p_2, q_2)$  be two arbitrary graphs. Then,

$$SO(H_1[H_2]_Q) \leq p_2^3(SO(L(H_1)) + M_1(H_1)) + p_1SO(H_2) + (2p_2)^{1/2} \left[ M_{1/2}(H_1)\chi_{1/2}(H_2) + p_2M_{3/2}(H_1)M_{1/2}(H_2) \right] + 2^{3/2}(2^{1/2} + 1)p_2q_1q_2 + p_2^2 \left[ 2\chi_{1/2}(H_1) + 2^{1/2}p_2 \left\{ M_1(H_1) - 2q_1 + 2^{1/2}\chi_{1/2}(L(H_1)) \right\} \right]$$

*Proof.*

$$\begin{aligned} SO(H_1[H_2]_Q) &= \sum_{(y_1, z_1)(y_2, z_2) \in E(H_1[H_2]_Q)} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\ &= \sum_{y_1=y_2 \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\ &\quad + \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{y_1 y_2 \in E(Q(H_1))} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\ &= Q_1 + Q_2 \end{aligned}$$

$$\begin{aligned} Q_1 &= \sum_{y_1=y_2 \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\ &= \sum_{y \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{(p_2 d_{Q(H_1)}(y) + d_{H_2}(z_1))^2 + (p_2 d_{Q(H_1)}(y) + d_{H_2}(z_2))^2} \\ &\leq p_1SO(H_2) + (2p_2)^{1/2}M_{1/2}(H_1)\chi_{1/2}(H_2) + 2^{3/2}p_2q_1q_2 \end{aligned}$$

$$\begin{aligned} Q_2 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{y_1 y_2 \in E(Q(H_1))} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\ &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(Q(H_1)) \\ y_1 \in V(H_1) \\ y_2 \in V(Q(H_1)) \setminus V(H_1)}} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\ &\quad + \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(Q(H_1)) \\ y_1, y_2 \in V(Q(H_1)) \setminus V(H_1)}} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\ &= Q'_2 + Q''_2 \end{aligned}$$



$$\begin{aligned}
 Q'_2 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(Q(H_1)) \\ y_1 \in V(H_1) \\ y_2 \in V(Q(H_1)) \setminus V(H_1)}} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\
 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(Q(H_1)) \\ y_1 \in V(H_1) \\ y_2 \in V(Q(H_1)) \setminus V(H_1)}} \sqrt{(p_2 d_{Q(H_1)}(y_1) + d_{H_2}(z_1))^2 + (p_2 d_{Q(H_1)}(y_2))^2} \\
 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(Q(H_1)) \\ y_1 \in V(H_1) \\ y_2 \in V(Q(H_1)) \setminus V(H_1)}} \sqrt{p_2^2 d_{H_1}(y_1)^2 + d_{H_2}(z_1)^2 + 2p_2 d_{H_1}(y_1) d_{H_2}(z_1) + p_2^2 d_{Q(H_1)}(y_2)^2} \\
 &\leq p_2^3 M_1(H_1) + (2q_2)p_2(2q_1) + (2p_2^3)^{1/2} M_{3/2}(H_1) M_{1/2}(H_2) + 2p_2^2 \chi_{1/2}(H_1)
 \end{aligned}$$

$$\begin{aligned}
 Q''_2 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(Q(H_1)) \\ y_1, y_2 \in V(Q(H_1)) \setminus V(H_1)}} \sqrt{d(y_1, z_1)^2 + d(y_2, z_2)^2} \\
 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(Q(H_1)) \\ y_1, y_2 \in V(Q(H_1)) \setminus V(H_1)}} \sqrt{(p_2 d_{Q(H_1)}(y_1))^2 + (p_2 d_{Q(H_1)}(y_2))^2} \\
 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{r_i, r_j, r_k \in E(H_1)} \sqrt{p_2^2 [(d_{H_1}(r_i) + d_{H_1}(r_j))^2 + (d_{H_1}(r_j) + d_{H_1}(r_k))^2]} \\
 &= p_2^3 \sum_{S_i, S_j \in E(L(H_1))} \sqrt{(d_{L(H_1)}(S_i) + 2)^2 + (d_{L(H_1)}(S_j) + 2)^2} \\
 &\leq p_2^3 [SO(L(H_1)) + 2^{3/2} (\frac{M_1(H_1)}{2} - q_1) + 2\chi_{1/2}(L(H_1))]
 \end{aligned}$$

From all the computations,

$$\begin{aligned}
 SO(H_1[H_2]_Q) &\leq p_2^3 (SO(L(H_1)) + M_1(H_1)) + p_1 SO(H_2) + (2p_2)^{1/2} [M_{1/2}(H_1) \chi_{1/2}(H_2) + p_2 M_{3/2}(H_1) \\
 &\quad M_{1/2}(H_2)] + 2^{3/2} (2^{1/2} + 1) p_2 q_1 q_2 + p_2^2 [2\chi_{1/2}(H_1) + 2^{1/2} p_2 \{M_1(H_1) - 2q_1 + 2^{1/2} \chi_{1/2}(L(H_1))\}]
 \end{aligned}$$

This concludes the result. □

**Theorem 2.4.** Assume  $H_1(p_1, q_1)$  and  $H_2(p_2, q_2)$  be two arbitrary graphs. Then,

$$\begin{aligned}
 SO(H_1[H_2]_T) &\leq 2p_2^3 (M_1(H_1) + SO(H_1)) + p_1 SO(H_2) + 2(p_2)^{1/2} [M_{1/2}(H_1) \chi_{1/2}(H_2) + p_2 M_{3/2}(H_1) \\
 &\quad M_{1/2}(H_2)] + p_2^2 [2\chi_{1/2}(H_1) + p_2 \{SO(L(H_1)) + 2^{1/2} (M_1(H_1) - 2q_1) + 2\chi_{1/2}(L(H_1))\}] \\
 &\quad + 4(2)^{1/2} (2^{1/2} + 1) p_2 q_1 q_2 + (8p_2^5 \Delta_{H_1} \Delta_{H_2})^{1/2} q_1
 \end{aligned}$$

Proof.

$$\begin{aligned}
 SO(H_1[H_2]_T) &= \sum_{(y_1, z_1)(y_2, z_2) \in E(H_1[H_2]_T)} \sqrt{d_{H_1[H_2]_T}(y_1, z_1)^2 + d_{H_1[H_2]_T}(y_2, z_2)^2} \\
 &= \sum_{y \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{d_{H_1[H_2]_T}(y, z_1)^2 + d_{H_1[H_2]_T}(y, z_2)^2} \\
 &+ \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{y_1 y_2 \in E(T(H_1))} \sqrt{d_{H_1[H_2]_T}(y_1, z_1)^2 + d_{H_1[H_2]_T}(y_2, z_2)^2} \\
 &= \sum_{y \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{d_{H_1[H_2]_R}(y, z_1)^2 + d_{H_1[H_2]_R}(y, z_2)^2} \\
 &+ \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{y_1 y_2 \in E(H_1)} \sqrt{d_{H_1[H_2]_R}(y_1, z_1)^2 + d_{H_1[H_2]_R}(y_2, z_2)^2} \\
 &+ \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(T(H_1)) \\ y_1 \in V(H_1) \\ y_2 \in V(T(H_1)) \setminus V(H_1)}} \sqrt{d_{H_1[H_2]_R}(y_1, z_1)^2 + d_{H_1[H_2]_Q}(y_2, z_2)^2} \\
 &+ \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(T(H_1)) \\ y_1, y_2 \in V(T(H_1)) \setminus V(H_1)}} \sqrt{d_{H_1[H_2]_Q}(y_1, z_1)^2 + d_{H_1[H_2]_Q}(y_2, z_2)^2} \\
 &= T_1 + T_2 + T_3 + T_4
 \end{aligned}$$

From the theorems (2.1),(2.2),(2.3), we have

$$\begin{aligned}
 T_1 &= \sum_{y \in V(H_1)} \sum_{z_1 z_2 \in E(H_2)} \sqrt{d_{H_1[H_2]_R}(y, z_1)^2 + d_{H_1[H_2]_R}(y, z_2)^2} \\
 &\leq p_1 SO(H_2) + 2(p_2)^{1/2} M_{1/2}(H_1) \chi_{1/2}(H_2) + 4(2)^{1/2} p_2 q_1 q_2 \\
 T_2 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{y_1 y_2 \in E(H_1)} \sqrt{d_{H_1[H_2]_R}(y_1, z_1)^2 + d_{H_1[H_2]_R}(y_2, z_2)^2} \\
 &\leq 2p_2^3 SO(H_1) + 4p_2 q_1 q_2 + (8p_2^5 \Delta_{H_1} \Delta_{H_2})^{1/2} q_1 \\
 T_3 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(T(H_1)) \\ y_1 \in V(H_1) \\ y_2 \in V(T(H_1)) \setminus V(H_1)}} \sqrt{d_{H_1[H_2]_R}(y_1, z_1)^2 + d_{H_1[H_2]_Q}(y_2, z_2)^2} \\
 &= \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(T(H_1)) \\ y_1 \in V(H_1) \\ y_2 \in V(T(H_1)) \setminus V(H_1)}} \sqrt{(2p_2 d_{H_1}(y_1) + d_{H_1}(z_2))^2 + (p_2 d_{Q(H_1)}(y_2))^2} \\
 &\leq 2p_2^2 (p_2 M_1(H_1) + \chi_{1/2}(H_1)) + 2p_2^{3/2} M_{1/2}(H_2) M_{3/2}(H_1) + 4p_2 q_1 q_2
 \end{aligned}$$

$$T_4 = \sum_{z_1 \in V(H_2)} \sum_{z_2 \in V(H_2)} \sum_{\substack{y_1 y_2 \in E(T(H_1)) \\ y_1, y_2 \in V(T(H_1)) \setminus V(H_1)}} \sqrt{d_{H_1[H_2]_Q}(y_1, z_1)^2 + d_{H_1[H_2]_Q}(y_2, z_2)^2}$$

$$\leq p_2^3 \left[ SO(L(H_1)) + 2^{1/2}(M_1(H_1) - 2q_1) + 2\chi_{1/2}(L(H_1)) \right]$$

Hence from all the computations,

$$SO(H_1[H_2]_T) \leq 2p_2^3(M_1(H_1) + SO(H_1)) + p_1 SO(H_2) + 2(p_2)^{1/2} [M_{1/2}(H_1)\chi_{1/2}(H_2) + p_2 M_{3/2}(H_1) M_{1/2}(H_2)] + p_2^2 [2\chi_{1/2}(H_1) + p_2 \{SO(L(H_1)) + 2^{1/2}(M_1(H_1) - 2q_1) + 2\chi_{1/2}(L(H_1))\}] + 4(2)^{1/2} (2^{1/2} + 1) p_2 q_1 q_2 + (8p_2^5 \Delta_{H_1} \Delta_{H_2})^{1/2} q_1$$

This concludes the result. □

**Corollary 2.5.** *The upper bound for the Sombor index of the lexicographic product versions of the path graphs is given as:*

- $SO(P_m[P_n]_S) \leq 8mn^3 + 6.83mn^2 - 10n^3 - 4mn - 6.83n^2 - 4.01m + 6.83n + n^{1/2} (5.66mn^2 + 0.69mn - 7.31n^2 - 5.07m + 1.94n + 2.97)$
- $SO(P_m[P_n]_R) \leq 17.66mn^3 + 13.66mn^2 - 24.03n^3 - 10.83mn - 13.66n^2 - 4.01m + 13.66n + n^{1/2} (8mn^2 + 0.97mn - 10.34n^2 - 7.17m + 2.75n + 4.20) + (32n^5)^{1/2}(m - 1)$
- $SO(P_m[P_n]_Q) \leq 13.66mn^3 + 10.83mn^2 - 27.57n^3 - 4mn - 11.90n^2 - 4.01m + 6.83n + n^{1/2} (5.66mn^2 + 0.69mn - 7.31n^2 - 5.07m + 1.94n + 2.97)$
- $SO(P_m[P_n]_T) \leq 23.31mn^3 + 17.66mn^2 - 41.60n^3 - 10.83mn - 18.73n^2 - 4.01m + 13.66n + n^{1/2} (8mn^2 + 0.97mn - 10.34n^2 - 7.17m + 2.75n + 4.20) + (32n^5)^{1/2}(m - 1)$

**Corollary 2.6.** *The upper bound for the Sombor index of the lexicographic product versions of the cycle graphs is given as:*

- $SO(C_m[C_n]_S) \leq 8mn^3 + 6.83mn^2 + 2.83mn + n^{1/2} (5.66mn^2 + 4mn)$
- $SO(C_m[C_n]_R) \leq 17.66mn^3 + 13.66mn^2 + 2.83mn + n^{1/2} (8mn^2 + 5.66) + (32n^5)^{1/2} m$
- $SO(C_m[C_n]_Q) \leq 13.66mn^3 + 10.83mn^2 + 2.83mn + n^{1/2} (5.66mn^2 + 4mn)$
- $SO(C_m[C_n]_T) \leq 23.31mn^3 + 17.66mn^2 + 2.83mn + n^{1/2} (8mn^2 + 5.66mn) + (32n^5)^{1/2} m$

**Corollary 2.7.** *The upper bound for the Sombor index of the lexicographic product versions involving path and cycle graphs is given as:*

- $SO(P_m[C_n]_S) \leq 8mn^3 + 6.83mn^2 - 10n^3 + 2.83mn - 6.83n^2 + n^{1/2} (5.66n^3 - 3.31n^2 - 2.34n)$
- $SO(P_m[C_n]_R) \leq 17.66mn^3 + 13.66mn^2 - 24.03n^3 + 2.83mn - 13.66n^2 + n^{1/2} (13.66mn^2 + 5.66mn - 15.997n^2 - 3.31n)$
- $SO(P_m[C_n]_Q) \leq 13.66mn^3 + 10.83mn^2 - 27.57n^3 + 2.83mn - 11.90n^2 + n^{1/2} (5.66mn^2 + 4mn - 7.31n^2 - 2.34n)$

- $SO(P_m[C_n]_T) \leq 23.31mn^3 + 17.66mn^2 - 41.60n^3 + 2.83mn - 18.73n^2 + n^{1/2} \left( 13.66mn^2 + 5.66mn - 16n^2 - 3.31n \right)$

**Corollary 2.8.** *The upper bound for the Sombor index of the lexicographic product versions involving cycle and path graphs is given as:*

- $SO(C_n[P_m]_S) \leq 8m^3n + 6.83m^2n - 4mn - 4.01n + m^{1/2} \left( 5.66m^2n + 0.69mn - 5.07n \right)$
- $SO(C_n[P_m]_R) \leq 17.66m^3n + 13.66m^2n - 10.83mn - 4.01n + m^{1/2} \left( 13.66m^2n + 0.97mn - 7.17n \right)$
- $SO(C_n[P_m]_Q) \leq 13.66m^3n + 10.83m^2n - 4mn - 4.01n + m^{1/2} \left( 5.66m^2n + 0.68mn - 5.07n \right)$
- $SO(C_n[P_m]_T) \leq 23.31m^3n + 17.65m^2n - 10.83mn - 4.01n + m^{1/2} \left( 13.66m^2n + 0.97mn - 7.17n \right)$

**Theorem 2.9.** *Assume  $H_1(p_1, q_1)$  and  $H_2(p_2, q_2)$  be two arbitrary graphs. Then,*

- $SO(H_1 \otimes_S H_2) < (M_1(H_1) + 4q_1)M_1(H_2)$
- $SO(H_1 \otimes_R H_2) < 4(M_1(H_1) + q_1)M_1(H_2)$
- $SO(H_1 \otimes_Q H_2) < (M_1(H_1) + HM_1(H_1))M_1(H_2)$
- $SO(H_1 \otimes_T H_2) < (4M_1(H_1) + HM_1(H_1))M_1(H_2)$

**Corollary 2.10.** *The upper bound for the Sombor index of the tensor product versions of the path graphs is given as:*

- $SO(P_m \otimes_S P_n) < 4(4m - 5)(2n - 3)$
- $SO(P_m \otimes_R P_n) < 8(5m - 7)(2n - 3)$
- $SO(P_m \otimes_Q P_n) < 8(5m - 9)(2n - 3)$
- $SO(P_m \otimes_T P_n) < 4(16m - 27)(2n - 3)$

**Corollary 2.11.** *The upper bound for the Sombor index of the tensor product versions of the cycle graphs is given as:*

- $SO(C_m \otimes_S C_n) < 32mn$
- $SO(C_m \otimes_R C_n) < 80mn$
- $SO(C_m \otimes_Q C_n) < 80mn$
- $SO(C_m \otimes_T C_n) < 128mn$

**2.2. Second Hyper Zagreb Index and First Reformulated Zagreb Index based on four operations related to the tensor product.** In this section, we determine the evaluated expressions of the second hyper-Zagreb index and the first reformulated Zagreb index of the tensor products based on the four graphs and further, we also obtain the expression values for the standard path and cycle graphs.

**Theorem 2.12.** *Assume  $H_1(p_1, q_1)$  and  $H_2(p_2, q_2)$  be two arbitrary graphs and  $F \in \{S, R, Q, T\}$ . Then,*

$$HM_2(H_1 \otimes_F H_2) = 2HM_2(F(H_1))HM_2(H_2)$$

Proof.

$$\begin{aligned}
 HM_2(H_1 \otimes_F H_2) &= \sum_{(y_1, z_1)(y_2, z_2) \in E(H_1 \otimes_F H_2)} (d_{H_1 \otimes_F H_2}(y_1, z_1) \cdot d_{H_1 \otimes_F H_2}(y_2, z_2))^2 \\
 &= 2 \sum_{y_1 y_2 \in E(F(H_1))} \sum_{z_1 z_2 \in E(H_2)} \left[ (d_{F(H_1)}(y_1) d_{H_2}(z_1)) \cdot (d_{F(H_1)}(y_2) d_{H_2}(z_2)) \right]^2 \\
 &= 2 \sum_{y_1 y_2 \in E(F(H_1))} (d_{F(H_1)}(y_1) \cdot d_{F(H_1)}(y_2))^2 \sum_{z_1 z_2 \in E(H_2)} (d_{H_2}(z_1) \cdot d_{H_2}(z_2))^2 \\
 &= 2HM_2(F(H_1))HM_2(H_2)
 \end{aligned}$$

□

**Corollary 2.13.** Assume  $H_1(p_1, q_1)$  and  $H_2(p_2, q_2)$  be two arbitrary graphs and  $F \in \{S, R, Q, T\}$ . Then,

- $HM_2(H_1 \otimes_S H_2) = 8F(H_1)HM_2(H_2)$
- $HM_2(H_1 \otimes_R H_2) = 32(HM_2(H_1) + F(H_1))HM_2(H_2)$
- $HM_2(H_1 \otimes_Q H_2) \leq 16\Delta_{H_1}^4 M_1(H_1)HM_2(H_2)$
- $HM_2(H_1 \otimes_T H_2) \leq 16\Delta_{H_1}^4 (M_1(H_1) + 4q_1)HM_2(H_2)$

**Theorem 2.14.** Assume  $H_1(p_1, q_1)$  and  $H_2(p_2, q_2)$  be two arbitrary graphs. Then,

- $EM_1(H_1 \otimes_S H_2) = (\mathbf{F}(H_1) + 8q_1)\mathbf{F}(H_2) + 8M_2(H_2)M_1(H_1) - 4M_1(H_2)(M_1(H_1) + 4q_1) + 16q_1q_2$
- $EM_1(H_1 \otimes_R H_2) = 8\mathbf{F}(H_2)(\mathbf{F}(H_1) + q_1) + 16M_2(H_2)(M_1(H_1) + M_2(H_1)) - 16(M_1(H_1) + q_1)M_1(H_2) + 24q_1q_2$
- $EM_1(H_1 \otimes_Q H_2) = \mathbf{F}(H_2)(\mathbf{F}(H_1) + \chi_3(H_1)) + 4q_2(M_1(H_1) + 2q_1) + 4(2EM_1(H_1) + EM_2(H_1) + 2\{M_1(H_1) + M_2(H_1)\} + \mathbf{F}(H_1) - 4q_1)M_2(H_2) - 4(M_1(H_1) + HM_1(H_1))M_1(H_2)$
- $EM_1(H_1 \otimes_T H_2) = \mathbf{F}(H_2)(8\mathbf{F}(H_1) + \chi_3(H_1)) + 4(2M_1(H_1) + 8M_2(H_1) + 2EM_1(H_1) + EM_2(H_1) + 2\mathbf{F}(H_1) - 4q_1)M_2(H_2) - 4(4M_1(H_1) + HM_1(H_1))M_1(H_2) + 4q_2(M_1(H_1) + 4q_1)$

**Corollary 2.15.** The exact expressions for the first reformulated Zagreb index for the path graphs  $P_m, P_n$  are:

- $EM_1(P_m \otimes_S P_n) = 144mn - 304m - 224n + 468$
- $EM_1(P_m \otimes_R P_n) = 792mn - 1576m - 1432n + 2824$
- $EM_1(P_m \otimes_Q P_n) = 792mn - 1576m - 1792n + 3536$
- $EM_1(P_m \otimes_T P_n) = 1568mn - 3104m - 3288n + 6468$

**Corollary 2.16.** The exact expressions for the first reformulated Zagreb index for the cycle graphs  $C_m, C_n$  are:

- $EM_1(C_m \otimes_S C_n) = 144mn$
- $EM_1(C_m \otimes_R C_n) = 792mn$
- $EM_1(C_m \otimes_Q C_n) = 792mn$

- $EM_1(C_m \otimes_T C_n) = 800mn$

**Corollary 2.17.** The exact expressions for the first reformulated Zagreb index of  $P_m \otimes_F C_n$  ( $F \in \{S, R, Q, T\}$ ) are:

- $EM_1(P_m \otimes_S C_n) = 144mn - 224n$
- $EM_1(P_m \otimes_R C_n) = 792mn - 1432n$
- $EM_1(P_m \otimes_Q C_n) = \begin{cases} 536mn + 512n^2 - 2336n, & m = 2 \\ 280mn + 512n^2 - 1792n, & m \geq 3 \end{cases}$
- $EM_1(P_m \otimes_T C_n) = \begin{cases} 1824mn - 3832n, & m = 2 \\ 1568mn - 3288n, & m \geq 3 \end{cases}$

**Corollary 2.18.** The exact expressions for the first reformulated Zagreb index of  $C_n \otimes_F P_m$  ( $F \in \{S, R, Q, T\}$ ) are:

- $EM_1(C_n \otimes_S P_m) = 144mn - 304n$
- $EM_1(C_n \otimes_R P_m) = 792mn - 1576n$
- $EM_1(C_n \otimes_Q P_m) = 792mn - 1576n$
- $EM_1(C_n \otimes_T P_m) = 1568mn - 3104n$

**2.3. Graphical Illustration.** The graphical depiction of the evaluated expression of the First Reformulated Zagreb index for the  $P_m \otimes_F P_n$  and  $C_m \otimes_F C_n$  graph operation is shown in Figure 4 where  $F \in \{S, R, Q, T\}$ .




FIGURE 4. Graphical illustration of  $EM_1(P_m \otimes_F P_n)$ (left) and  $EM_1(C_m \otimes_F C_n)$ (right)

The graphical depiction of the evaluated expression of the First Reformulated Zagreb index for the  $P_m \otimes_F C_n$  and  $C_n \otimes_F P_m$  graph operation is shown in Figure (5) where  $F \in \{S, R, Q, T\}$ .




FIGURE 5. Graphical illustration of exact expression of  $EM_1(P_m \otimes_F C_n)$ (left) and  $EM_1(C_n \otimes_F P_m)$ (right)

From the figures 4 and 5, we can assert that the first reformulated Zagreb index for the tensor product graph operations involving the four derived graphs holds the greatest value for total graph product version in all the cases of path and cycle graphs.

### 3. Conclusion

Molecular descriptors have a considerable influence in the comprehension of medical techniques and social network theory. In this paper, we execute the determination of the upper bound expressions of the Sombor index of the lexicographic and tensor product of two graphs based on the four operations involving the subdivision graphs, semi-total point graphs, semi-total line graphs and the total graphs.

The formulation of the exact expressions for the second hyper Zagreb index and the first reformulated Zagreb index of the tensor products of the graphs based on the four graphical operations is carried out. Further, the descriptor bounds and values with respect to standard path and cycle graphs are obtained and for the proper illustration of the notion, the graphical comparison for the first reformulated Zagreb index has been depicted in the paper.

For the future scope of the study, one can look into the formulation of the indices for the generalized  $F$ -lexicographic and  $F$ -tensor product versions where  $F \in \{S, R, Q, T\}$ . The formulations of the topological index of the complex network structures which are of the forms of the resultant path and cycle graph lexicographic and tensor product graph combinations is obtained just with the knowledge of the base graph topological index computations. The results have strong significance towards wide and stringent investigation of bond-additive indices and graph transformed structural networks.

### Acknowledgments

The authors wish to thank the anonymous referees for their valuable suggestions to improvise the manuscript.

### REFERENCES

- [1] S. Akhter, R. Farooq and S. Pirzada, Exact formulae of general sum-connectivity index for some graph operations, *Mat. Vesnik*, **70** no. 3 (2018) 267–282.
- [2] N. E. Arif, A. H. Karim and R. Hasni, Sombor index of some graph operations, *Int. J. Nonlinear Anal. Appl.*, **13** no. 1 (2022) 2561–2571.
- [3] B. Basavanagoud and A. P. Barangi,  $F$ -index and hyper-Zagreb index of four new tensor products of graphs and their complements, *Discrete Math. Algorithms Appl.*, **11** no. 03 (2019) 14 pp.
- [4] B. Basavanagoud, V. Desai, K. G. Mirajkar, B. Pooja and I. N. Cangul, Four new tensor products of graphs and their Zagreb indices and coindices, *Electron. J. Math. Anal. Appl.*, **8** no. 1 (2020) 209–219.
- [5] B. Bommanahal, M. Sayyed and B. Pooja, Topological indices of four new tensor products of graphs, *Annals of Mathematics and Computer Science*, **9** (2022) 52–66.
- [6] K. C. Das, A. S. Çevik, I. N. Cangul and Y. Shang, On Sombor index, *Symmetry*, **13** no. 1 (2021) Article number: 140.
- [7] N. De,  $F$ -index of graphs based on four operations related to the lexicographic product, *Malaya J. Mat.*, **8** no. 2 (2020) 397–404.
- [8] N. De, S. M. A. Nayeem and A. Pal,  $F$ -index of some graph operations, *Discrete Math. Algorithms Appl.*, **8** no. 2 (2016) 17 pp.
- [9] N. Dehgardi, S. Sheikholeslami and M. Soroudi, Some distance based indices of graphs based on four new operations related to the lexicographic product, *Carpathian Math. Publ.*, **11** no. 2 (2019) 258–267.
- [10] M. Eliasi and B. Taeri, Four new sums of graphs and their Wiener indices, *Discrete Appl. Math.*, **157** no. 4 (2009) 794–803.
- [11] M. Farahani, M. R. Kanna and R. P. Kumar, On the hyper-Zagreb indices of nano-structures, *Asian Academic Research Journal of Multidisciplinary*, **3** no. 1 (2016) 115–123.
- [12] B. Furtula and I. Gutman, A forgotten topological index, *J. Math. Chem.*, **53** no. 4 (2015) 1184–1190.
- [13] W. Gao, M. R. Farahani, M. K. Siddiqui, and M. K. Jamil, On the first and second Zagreb and first and second hyper-Zagreb indices of carbon nanocones  $CNC_k[n]$ , *J. Comput. Theor. Nanosci.*, **13** no. 10 (2016) 7475–7482.
- [14] I. Gutman, Geometric approach to degree-based topological indices: Sombor indices, *MATCH Commun. Math. Comput. Chem.*, **86** no. 1 (2021) 11–16.
- [15] I. Gutman, Some basic properties of Sombor indices, *Open J. Discrete Appl. Math.*, **4** no. 1 (2021) 1–3.
- [16] I. Gutman, V. Kulli, B. Chaluvvaraju, and H. Boregowda, On Banhatti and Zagreb indices, *J. Int. Math. Virtual Inst.*, **7** no. 1 (2017) 53–67.
- [17] I. Gutman, E. Milovanović, and I. Milovanović, Beyond the Zagreb indices, *AKCE Int. J. Graphs Comb.*, **17** (2020) 74–85.



- [18] I. Gutman and N. Trinajstić, Graph theory and molecular orbitals. Total  $\varphi$ -electron energy of alternant hydrocarbons, *Chem. Phys. Lett.*, **17** no. 4 (1972) 535–538.
- [19] N. Harish, B. Sarveshkumar, and B. Chaluvaram, The reformulated Sombor index of a graph, *Trans. Comb.*, **13** no. 1 (2024) 1–16.
- [20] Harisha, P. S. Ranjini, V. Lokesh, K-Banhatti indices for special graphs and vertex gluing graphs, *International J. Math. Combin.*, **4** (2020) 89–99.
- [21] B. Horoldagva and X. Chunlei, On Sombor index of graphs, *MATCH Commun. Math. Comput. Chem.*, **86** no. 3 (2021) 703–713.
- [22] W. Imrich and S. Klavžar, *Product graphs. Structure and recognition*, Wiley-Interscience Series in Discrete Mathematics and Optimization, Wiley-Interscience, New York, 2000.
- [23] V. Kulli, Sombor indices of certain graph operators, *Int. J. Engin. Sci. Res. Technol.*, **10** no. 1 (2021) 127–134.
- [24] V. R. Kulli, N. Harish, B. Chaluvaram, and I. Gutman, Mathematical properties of KG Sombor index, *Bull. Int. Math. Virtual Inst.*, **12** no. 2 (2022) 379–386.
- [25] X. Li and J. Zheng, A unified approach to the extremal trees for different indices, *MATCH Commun. Math. Comput. Chem.*, **54** no. 1 (2005) 195–208.
- [26] H. Liu, L. You, Y. Huang and Z. Tang, On extremal Sombor indices of chemical graphs, and beyond, *MATCH Commun. Math. Comput. Chem.*, **89** (2023) 415–436.
- [27] V. Lokesh, S. Jain, A. Cevik, and I. Cangul, New results on the  $F$ -index of graphs based on corona-type products of graphs, In *Proc. Jangjeon Math. Soc.*, **23** (2020) 141–148.
- [28] V. Lokesh, K. Z. Yasmeen, M. Manjunath and M. Venkatachalam, Lower and upper bounds of SK indices of  $F$ -join of graphs, *Int. J. Open Problems Compt. Math.*, **13** no. 4 (2020) 13–25.
- [29] A. Miličević, S. Nikolić and N. Trinajstić, On reformulated Zagreb indices, *Mol. Divers.*, **8** no. 4 (2004) 393–399.
- [30] M. Nadjafi-Arani and H. Khodashenas, Distance-based topological indices of tensor product of graphs, *Iranian J. Math. Chem.*, **3** no. 1 (2012) 45–53.
- [31] K. Pattabiraman, Four new operations related to composition and their hyper-Zagreb index, *Southeast Asian Bull. Math.*, **43** no. 6 (2019).
- [32] E. Sampathkumar and S. Chikkodimath, Semitotal graphs of a graph-I, *J. Karnatak Univ. Sci.*, **18** (1973) 274–280.
- [33] D. Sarala, H. Deng, S. Ayyaswamy, and S. Balachandran, The Zagreb indices of graphs based on four new operations related to the lexicographic product, *Appl. Math. Comput.*, **309** (2017) 156–169.
- [34] G. Shirdel, H. Rezapour and A. Sayadi, The hyper-Zagreb index of graph operations, *Iranian J. Math. Chem.*, **4** no. 2 (2013) 213–220.
- [35] B. Shwetha Shetty, V. Lokesh and P. Ranjini, On the harmonic index of graph operations, *Trans. comb.*, **4** no. 4 (2015) 5–14.
- [36] H. M. A. Siddiqui, S. Baby, M. F. Nadeem and M. K. Shafiq, Bounds of some degree based indices of lexicographic product of some connected graphs, *Polycyclic Aromat. Compd.*, **42** no. 5 (2022) 2568–2580.
- [37] Z. Wang, Y. Mao, Y. Li and B. Furtula, On relations between Sombor and other degree-based indices, *J. Appl. Math. Comput.*, **68** (2022) 1–17.
- [38] D. B. West, *Introduction to graph theory*, Prentice Hall, Inc., Upper Saddle River, NJ, 1996.

- [39] Z. Yarahmadi, S. Moradi and T. Došlić, Eccentric connectivity index of graphs with subdivided edges, *Electron. Notes Discrete Math.*, **45** (2014) 167–176.
- [40] B. Zhou and N. Trinajstić, On a novel connectivity index, *J. Math. Chem.*, **46** no. 4 (2009) 1252–1270.
- [41] B. Zhou and N. Trinajstić, On general sum-connectivity index, *J. Math. Chem.*, **47** no. 1 (2010) 210–218.

**Ishita Sarkar**

Department of Mathematics, CHRIST (Deemed to be University), Bengaluru 560029, India

Email: [ishita.sarkar@res.christuniversity.in](mailto:ishita.sarkar@res.christuniversity.in)

**Manjunath Nanjappa**

Department of Mathematics, CHRIST (Deemed to be University), Bengaluru 560029, India

Email: [manjunath.nanjappa@christuniversity.in](mailto:manjunath.nanjappa@christuniversity.in)